

共同演者論文

1. Yamaji K, Shiomi H, Morimoto T, Nakatsuma K, Toyota T, Ono K, Furukawa Y, Nakagawa Y, Kadota K, Ando K, Shirai S, Onodera T, Watanabe H, Natsuaki M, Sakata R, Hanyu M, Nishiwaki N, Komiya T, Kimura T. Effects of Age and Sex on Clinical Outcomes after Percutaneous Coronary Intervention Relative to Coronary Artery Bypass Grafting in Patients with Triple Vessel Coronary Artery Disease. *Circulation*. 2016 Mar 23. pii: CIRCULATIONAHA.115.020955. [Epub ahead of print]
2. Wöhrle J, Markovic S, Rottbauer W, Muramatsu T, Kadota K, Vázquez-González N, Odenstedt J, Serra A, Antonucci D, Varenne O, Saito S, Wijns W. Bioresorbable polymer sirolimus-eluting coronary stent compared with permanent polymer everolimus-eluting coronary stent implantation for treatment of small vessel coronary artery disease: CENTURY II trial. *EuroIntervention*. 2016 Jun 12;12(2):e167-74. doi: 10.4244/EIJV12I2A30.
3. Nakatsuma K, Shiomi H, Morimoto T, Furukawa Y, Nakagawa Y, Ando K, Kadota K, Yamamoto T, Suwa S, Horie M, Kimura T; CREDO-Kyoto AMI investigators. Inter-Facility Transfer vs. Direct Admission of Patients With ST-Segment Elevation Acute Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. *Circ J*. 2016 Jun 28. [Epub ahead of print]
4. Kim MH, Mitsudo K, Jin CD, Kim TH, Cho YR, Park JS, Park K, Park TH, Serebruany V. Long-term clinical outcomes after successful and failed recanalization to native chronic Total occlusion: Insights from the Busan chronic Total occlusion (B-CTO) Registry. *Cardiovasc Revasc Med*. 2016 Jun;17(4):229-32.
5. Nakatsuma K, Shiomi H, Morimoto T, Ando K, Kadota K, Watanabe H, Taniguchi T, Yamamoto T, Furukawa Y, Nakagawa Y, Horie M, Kimura T; CREDO-Kyoto AMI investigators. Intravascular Ultrasound Guidance vs. Angiographic Guidance in Primary Percutaneous Coronary Intervention for ST-Segment Elevation Myocardial Infarction - Long-Term Clinical Outcomes From the CREDO-Kyoto AMI Registry. *Circ J*. 2016 Jan 25;80(2):477-84.
6. Taniguchi T, Morimoto T, Shiomi H, Ando K, Kanamori N, Murata K, Kitai T, Kawase Y, Izumi C, Miyake M, Mitsuoka H, Kato M, Hirano Y, Matsuda S, Nagao K, Inada T, Murakami T, Takeuchi Y, Yamane K, Toyofuku M, Ishii M, Minamino-Muta E, Kato T, Inoko M, Ikeda T,

- Komasa A, Ishii K, Hotta K, Higashitani N, Kato Y, Inuzuka Y, Maeda C, Jinnai T, Morikami Y, Sakata R, Kimura T; CURRENT AS Registry Investigators. Initial Surgical Versus Conservative Strategies in Patients With Asymptomatic Severe Aortic Stenosis. *J Am Coll Cardiol*. 2015 Dec 29;66(25):2827-38. doi: 10.1016/j.jacc.2015.10.001. Epub 2015 Oct 15.
7. Kimura T, Kozuma K, Tanabe K, Nakamura S, Yamane M, Muramatsu T, Saito S, Yajima J, Hagiwara N, Mitsudo K, Popma JJ, Serruys PW, Onuma Y, Ying S, Cao S, Staehr P, Cheong WF, Kusano H, Stone GW; ABSORB Japan Investigators. A randomized trial evaluating everolimus-eluting Absorb bioresorbable scaffolds vs. everolimus-eluting metallic stents in patients with coronary artery disease: ABSORB Japan. *Eur Heart J*. 2015 Dec 14;36(47):3332-42.
 8. Natsuaki M, Morimoto T, Yamamoto E, Shiomi H, Furukawa Y, Abe M, Nakao K, Ishikawa T, Kawai K, Yunoki K, Shimizu S, Akao M, Miki S, Yamamoto M, Okada H, Hoshino K, Kadota K, Morino Y, Igarashi K, Tanabe K, Kozuma K, Kimura T. One-year outcome of a prospective trial stopping dual antiplatelet therapy at 3 months after everolimus-eluting cobalt-chromium stent implantation: ShortT and Optimal duration of Dual AntiPlatelet Therapy after everolimus-eluting cobalt-chromium stent (STOPDAPT) trial. *Cardiovasc Interv Ther*. 2015 Oct 30. [Epub ahead of print]
 9. Yano M, Natsuaki M, Morimoto T, Nakagawa Y, Kawai K, Miyazaki S, Muramatsu T, Shiode N, Namura M, Sone T, Oshima S, Nishikawa H, Hiasa Y, Hayashi Y, Nobuyoshi M, Mitsudo K, Kimura T; j-Cypher Registry Investigators. Antiplatelet therapy discontinuation and stent thrombosis after sirolimus-eluting stent implantation: five-year outcome of the j-Cypher Registry. *Int J Cardiol*. 2015 Nov 15;199:296-301.
 10. Khand A, Patel B, Palmer N, Jones J, Andron M, Perry R, Mehrotra S, Mitsudo K. Retrograde Wiring of Collateral Channels of the Heart in Chronic Total Occlusions: A Systematic Review and Meta-Analysis of Safety, Feasibility, and Incremental Value in Achieving Revascularization. *Angiology*. 2015 Nov;66(10):925-32.
 11. Marui A, Kimura T, Nishiwaki N, Mitsudo K, Komiya T, Hanyu M, Shiomi H, Tanaka S, Sakata R; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Five-year outcomes of percutaneous versus surgical coronary revascularization in patients with diabetes mellitus (from the CREDO-Kyoto PCI/CABG Registry Cohort-2). *Am J Cardiol*. 2015 Apr 15;118(8):1063-72.

12. Natsuaki M, Kozuma K, Morimoto T, Kadota K, Muramatsu T, Nakagawa Y, Akasaka T, Igarashi K, Tanabe K, Morino Y, Ishikawa T, Nishikawa H, Awata M, Abe M, Okada H, Takatsu Y, Ogata N, Kimura K, Urasawa K, Tarutani Y, Shiode N, Kimura T. Final 3-Year Outcome of a Randomized Trial Comparing Second-Generation Drug-Eluting Stents Using Either Biodegradable Polymer or Durable Polymer: NOBORI Biolimus-Eluting Versus XIENCE/PROMUS Everolimus-Eluting Stent Trial. *Circ Cardiovasc Interv.* 2015;8(10):e002817.
13. Yamaji K, Shiomi H, Morimoto T, Toyota T, Ono K, Furukawa Y, Nakagawa Y, Kadota K, Ando K, Shirai S, Kato M, Takatsu Y, Doi O, Kambara H, Suwa S, Onodera T, Watanabe H, Natsuaki M, Kimura T. Influence of Sex on Long-Term Outcomes After Implantation of Bare-Metal Stent: A Multicenter Report From the Coronary Revascularization Demonstrating Outcome Study-Kyoto (CREDO-Kyoto) Registry Cohort-1. *Circulation.* 2015;132(24):2323-33.
14. Kubo T, Akasaka T, Kozuma K, Kimura K, Kawamura M, Sumiyoshi T, Ino Y, Morino Y, Tanabe K, Kadota K, Kimura T; RESET Investigators. Comparison of neointimal coverage between everolimus-eluting stents and sirolimus-eluting stents: an optical coherence tomography substudy of the RESET (Randomized Evaluation of Sirolimus-eluting versus Everolimus-eluting stent Trial). *EuroIntervention.* 2015;11(5):564-71.
15. Shimizu A, Mitsuhashi T, Nitta T, Mitamura H, Kurita T, Abe H, Nakazato Y, Sumitomo N, Kadota K, Kimura K, Okumura K; Committee for Implantable Devices Enrollment and Assessment in the Japanese Heart Rhythm Society. Japan Implantable Devices in Coronary Artery Disease (JID-CAD) study design. *J Arrhythm.* 2015;31(2):83-7.
16. Watanabe H, Shiomi H, Nakatsuma K, Morimoto T, Taniguchi T, Furukawa Y, Nakagawa Y, Horie M, Kimura T; CREDO - Kyoto AMI investigators, Kimura T, Sakata R, Marui A, Matsuda M, Mitsuoka H, Onoe M, Nakagawa Y, Yamanaka K, Fujiwara H, Takatsu Y, Ohno N, Nohara R, Murakami T, Takeda T, Nobuyoshi M, Iwabuchi M, Hanyu M, Tatami R, Matsushita T, Shirotani M, Nishiwaki N, Kita T, Furukawa Y, Okada Y, Kato H, Eizawa H, Is K, Tanaka M, Nakayama S, Lee JD, Nakano A, Koshiji T, Morioka K, Takizawa A, Shimamoto M, Yamazaki F, Takahashi M, Nishizawa J, Horie M, Takashima H, Tamura T, Aota M, Takahashi M, Tabata T, Tei C, Hamasaki S, Imoto Y, Yamamoto H, Kambara H, Doi O, Matsuda K, Nara M, Mitsudo K, Kadota K, Komiya T, Miki S, Mizoguchi T, Nakajima H, Ogawa H, Sugiyama S, Kawasaki

M, Moriyama S, Hattori R, Aoyama T, Araki M, Suwa S, Tanbara K, Kitagawa K, Yamauchi M, Okamoto N, Fujino Y, Tezuka S, Saeki A, Hanazawa M, Sato Y, Hibi C, Sasae H, Takinami E, Uchida Y, Yamamoto Y, Nishida S, Yoshimoto M, Maeda S, Miki I, Minematsu S, Abe M, Shiomi H, Tada T, Tazaki J, Kato Y, Hayano M, Tokushige A, Natsuaki M, Nakajima T. Clinical efficacy of thrombus aspiration on 5-year clinical outcomes in patients with ST-segment elevation acute myocardial infarction undergoing percutaneous coronary intervention. *J Am Heart Assoc.* 2015;4(6):e001962.

17. Watanabe Y, Hayashida K, Takayama M, Mitsudo K, Nanto S, Takanashi S, Komiya T, Kuratani T, Tobaru T, Goto T, Lefèvre T, Sawa Y, Morice MC. First direct comparison of clinical outcomes between European and Asian cohorts in transcatheter aortic valve implantation: the Massy study group vs. the PREVAIL JAPAN trial. *J Cardiol.* 2015 Feb;65(2):112-6.
18. Onuma Y, Kimura T, Räber L, Magro M, Girasis C, van Domburg R, Windecker S, Mitsudo K, Serruys PW; Bern-Rotterdam and j-Cypher registries. Differences in coronary risk factors, procedural characteristics, mortality and stent thrombosis between two all-comers percutaneous coronary intervention registries from Europe and Japan: a patient-level data analysis of the Bern-Rotterdam and j-Cypher registries. *EuroIntervention.* 2015 Sep;11(5):533-40.
19. Sawa Y, Takayama M, Mitsudo K, Nanto S, Takanashi S, Komiya T, Kuratani T, Tobaru T, Goto T. Clinical efficacy of transcatheter aortic valve replacement for severe aortic stenosis in high-risk patients: the PREVAIL JAPAN trial. *Surg Today.* 2015 Jan;45(1):34-43.
20. Shimamura K, Kubo T, Akasaka T, Kozuma K, Kimura K, Kawamura M, Sumiyoshi T, Ino Y, Yoshiyama M, Sonoda S, Igarashi K, Miyazawa A, Uzui H, Sakanoue Y, Shinke T, Morino Y, Tanabe K, Kadota K, Kimura T. Outcomes of everolimus-eluting stent incomplete stent apposition: a serial optical coherence tomography analysis. *Eur Heart J Cardiovasc Imaging.* 2015;16(1):23-8.
21. Yamaji K, Natsuaki M, Morimoto T, Ono K, Furukawa Y, Nakagawa Y, Kadota K, Ando K, Shirai S, Watanabe H, Shiomi H, Kimura T. Long-Term Outcomes After Coronary Stent Implantation in Patients Presenting With Versus Without Acute Myocardial Infarction (an Observation from Coronary Revascularization Demonstrating Outcome Study-Kyoto Registry Cohort-2). *Am J Cardiol.* 2015;116(1):15-23.

22. Komiya T, Ueno G, Kadota K, Mitsudo K, Okabayashi H, Nishiwaki N, Hanyu M, Kimura T, Tanaka S, Marui A, Sakata R; CREDO-Kyoto Investigators. An optimal strategy for coronary revascularization in patients with severe renal dysfunction. *Eur J Cardiothorac Surg.* 2015;48(2):293-300.
23. Watanabe H, Morimoto T, Natsuaki M, Furukawa Y, Nakagawa Y, Kadota K, Yamaji K, Ando K, Shizuta S, Shiomi H, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Tamura T, Shirotani M, Miki S, Matsuda M, Takahashi M, Ishii K, Tanaka M, Aoyama T, Doi O, Hattori R, Kato M, Suwa S, Takizawa A, Takatsu Y, Shinoda E, Eizawa H, Takeda T, Lee JD, Inoko M, Ogawa H, Hamasaki S, Horie M, Nohara R, Kambara H, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T, Kastrati A, Kimura T; CREDO-Kyoto PCI/CABG registry cohort-2 investigators. Antiplatelet therapy discontinuation and the risk of serious cardiovascular events after coronary stenting: observations from the CREDO-Kyoto Registry Cohort-2. *PLoS One.* 2015;10(4):e0124314.
24. Yamaji K, Kubo S, Inoue K, Kadota K, Kuramitsu S, Shirai S, Ando K, Nobuyoshi M, Mitsudo K, Kimura T. Association of localized hypersensitivity and in-stent neoatherosclerosis with the very late drug-eluting stent thrombosis. *PLoS One.* 2014;9(11):e113870.
25. Tazaki J, Morimoto T, Sakata R, Okabayashi H, Yamazaki F, Nishiwaki N, Mitsudo K, Kimura T; CREDO-Kyoto PCI/CABG registry cohort-2 investigators. Impact of statin therapy on patients with coronary heart disease and aortic aneurysm or dissection. *J Vasc Surg.* 2014 Sep;60(3):604-12.e2. doi:10.1016/j.jvs.2014.03.280. Epub 2014 May 3.
26. Ko E, Natsuaki M, Toyofuku M, Morimoto T, Matsumura Y, Oi M, Motohashi Y, Takahashi K, Kawase Y, Tanaka M, Kitada M, Yuzuki Y, Tamura T, Inoue K, Mitsudo K, Kimura T. Sirolimus-eluting stent implantation for ostial right coronary artery lesions: five-year outcomes from the j-Cypher registry. *Cardiovasc Interv Ther.* 2014 Jul;29(3):200-8.
27. Marui A, Kimura T, Nishiwaki N, Mitsudo K, Komiya T, Hanyu M, Shiomi H, Tanaka S, Sakata R; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Comparison of five-year outcomes of coronary artery bypass grafting versus percutaneous coronary intervention in patients with left ventricular ejection fractions $\leq 50\%$ versus $>50\%$ (from the CREDO-Kyoto PCI/CABG Registry Cohort-2). *Am J Cardiol.* 2014 Oct 1;114(7):988-96.
28. Taniguchi T, Shiomi H, Toyota T, Morimoto T, Akao M, Nakatsuma K, Ono K, Makiyama T, Shizuta S, Furukawa Y, Nakagawa Y, Ando K, Kadota K, Horie M, Kimura T. Effect of

preinfarction angina pectoris on long-term survival in patients with ST-segment elevation myocardial infarction who underwent primary percutaneous coronary intervention. *Am J Cardiol.* 2014 Oct 15;114(8):1179-86.

29. Kubo T, Akasaka T, Kozuma K, Kimura K, Fusazaki T, Okura H, Shinke T, Ino Y, Hasegawa T, Takashima H, Takamisawa I, Yamaguchi H, Igarashi K, Kadota K, Tanabe K, Nakagawa Y, Muramatsu T, Morino Y, Kimura T; NEXT Investigators. Vascular response to drug-eluting stent with biodegradable vs. durable polymer. Optical coherence tomography substudy of the NEXT. *Circ J.* 2014;78(10):2408-14.
30. Marui A, Kimura T, Nishiwaki N, Mitsudo K, Komiya T, Hanyu M, Shiomi H, Tanaka S, Sakata R; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Percutaneous coronary intervention versus coronary artery bypass grafting in patients with end-stage renal disease requiring dialysis (5-year outcomes of the CREDO-Kyoto PCI/CABG Registry Cohort-2). *Am J Cardiol.* 2014 Aug 15;114(4):555-61.
31. Tokushige A, Shiomi H, Morimoto T, Ono K, Furukawa Y, Nakagawa Y, Kadota K, Ando K, Shizuta S, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Hamasaki S, Ohishi M, Nakashima H, Mitsudo K, Nobuyoshi M, Kita T, Imoto Y, Sakata R, Okabayashi H, Hanyu M, Shimamoto M, Nishiwaki N, Komiya T, Kimura T; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Incidence and outcome of surgical procedures after coronary artery bypass grafting compared with those after percutaneous coronary intervention: a report from the Coronary Revascularization Demonstrating Outcome Study in Kyoto PCI/CABG Registry Cohort-2. *Circ Cardiovasc Interv.* 2014;7(4):482-91.
32. Campos CM, van Klaveren D, Iqbal J, Onuma Y, Zhang YJ, Garcia-Garcia HM, Morel MA, Farooq V, Shiomi H, Furukawa Y, Nakagawa Y, Kadota K, Lemos PA, Kimura T, Steyerberg EW, Serruys PW. Predictive Performance of SYNTAX Score II in Patients With Left Main and Multivessel Coronary Artery Disease-analysis of CREDO-Kyoto registry. *Circ J.* 2014;78(8):1942-9.
33. Goto K, Nakai K, Shizuta S, Morimoto T, Shiomi H, Natsuaki M, Yahata M, Ota C, Ono K, Makiyama T, Nakagawa Y, Furukawa Y, Kadota K, Takatsu Y, Tamura T, Takizawa A, Inada T, Doi O, Nohara R, Matsuda M, Takeda T, Kato M, Shirohani M, Eizawa H, Ishii K, Lee JD, Takahashi M, Horie M, Takahashi M, Miki S, Aoyama T, Suwa S, Hamasaki S, Ogawa H, Mitsudo K, Nobuyoshi M, Kita T, Kimura T; CREDO-Kyoto Registry Cohort-2 Investigators.

Anticoagulant and antiplatelet therapy in patients with atrial fibrillation undergoing percutaneous coronary intervention. *Am J Cardiol.* 2014;114(1):70-8.

34. Shiomi H, Kozuma K, Morimoto T, Igarashi K, Kadota K, Tanabe K, Morino Y, Akasaka T, Abe M, Suwa S, Muramatsu T, Kobayashi M, Dai K, Nakao K, Uematsu M, Tarutani Y, Fujii K, Simonton CA, Kimura T; RESET Investigators. Long-term clinical outcomes after everolimus- and sirolimus-eluting coronary stent implantation: final 3-year follow-up of the Randomized Evaluation of Sirolimus-Eluting Versus Everolimus-Eluting Stent Trial. *Circ Cardiovasc Interv.* 2014;7(3):343-54.
35. Shiomi H, Morimoto T, Makiyama T, Ono K, Furukawa Y, Nakagawa Y, Kadota K, Onodera T, Takatsu Y, Mitsudo K, Kita T, Sakata R, Okabayashi H, Hanyu M, Komiya T, Yamazaki F, Nishiwaki N, Kimura T; CREDO-Kyoto Investigators. Evolution in practice patterns and long-term outcomes of coronary revascularization from bare-metal stent era to drug-eluting stent era in Japan. *Am J Cardiol.* 2014;113(10):1652-9.
36. Kokunai Y, Nakata T, Furuta M, Sakata S, Kimura H, Aiba T, Yoshinaga M, Osaki Y, Nakamori M, Itoh H, Sato T, Kubota T, Kadota K, Shindo K, Mochizuki H, Shimizu W, Horie M, Okamura Y, Ohno K, Takahashi MP. A Kir3.4 mutation causes Andersen-Tawil syndrome by an inhibitory effect on Kir2.1. *Neurology.* 2014;82(12):1058-64.
37. Natsuaki M, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Yamaji K, Ando K, Shizuta S, Shiomi H, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Tamura T, Shirotani M, Miki S, Matsuda M, Takahashi M, Ishii K, Tanaka M, Aoyama T, Doi O, Hattori R, Kato M, Suwa S, Takizawa A, Takatsu Y, Shinoda E, Eizawa H, Takeda T, Lee JD, Inoko M, Ogawa H, Hamasaki S, Horie M, Nohara R, Kambara H, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T, Kimura T; CREDO-Kyoto PCI/CABG registry cohort-2 investigators. Late adverse events after implantation of sirolimus-eluting stent and bare-metal stent: long-term (5-7 years) follow-up of the Coronary Revascularization Demonstrating Outcome study-Kyoto registry Cohort-2. *Circ Cardiovasc Interv.* 2014;7(2):168-79.
38. Imai M, Kimura T, Morimoto T, Saito N, Shiomi H, Kawaguchi R, Kan H, Mukawa H, Fujita H, Ishise T, Hayashi F, Nagao K, Take S, Taniguchi H, Sakamoto H, Yamane T, Shiota K, Tamekiyo H, Okamura T, Kishi K, Miyazaki S, Yamamoto S, Yamaji K, Kawasaki T, Taguchi E, Nakajima H, Kosedo I, Tada T, Kadota K, Mitsudo K. Impact of angiographic peri-stent contrast staining (PSS) on late adverse events after sirolimus-eluting stent implantation: an

observation from the multicenter j-Cypher registry PSS substudy. *Cardiovasc Interv Ther.* 2014;29(3):226-36.

39. Kitai Y, Ozasa N, Morimoto T, Bao B, Furukawa Y, Nakagawa Y, Kadota K, Yanagita M, Shizuta S, Kimura T; CREDO-Kyoto registry investigators. Prognostic implications of anemia with or without chronic kidney disease in patients undergoing elective percutaneous coronary intervention. *Int J Cardiol.* 2013;168(6):5221-8.
40. Toyofuku M, Kimura T, Morimoto T, Hayashi Y, Shiode N, Nishikawa H, Nakao K, Shirota K, Kawai K, Hiasa Y, Kadota K, Nozaki Y, Isshiki T, Sone T, Mitsudo K; j-Cypher Registry Investigators. Comparison of 5-year outcomes in patients with and without unprotected left main coronary artery disease after treatment with sirolimus-eluting stents: insights from the j-Cypher registry. *JACC Cardiovasc Interv.* 2013;6(7):654-63.
41. Natsuaki M, Kozuma K, Morimoto T, Kadota K, Muramatsu T, Nakagawa Y, Akasaka T, Igarashi K, Tanabe K, Morino Y, Ishikawa T, Nishikawa H, Awata M, Abe M, Okada H, Takatsu Y, Ogata N, Kimura K, Urasawa K, Tarutani Y, Shiode N, Kimura T; NEXT Investigators. Biodegradable polymer biolimus-eluting stent versus durable polymer everolimus-eluting stent: a randomized, controlled, noninferiority trial. *J Am Coll Cardiol.* 2013;62(3):181-90.
42. Kozuma K, Kimura T, Kadota K, Suwa S, Kimura K, Iwabuchi M, Kawai K, Miyazawa A, Kawamura M, Nakao K, Asano R, Yamamoto T, Suzuki N, Aoki J, Kyono H, Nakazawa G, Tanabe K, Morino Y, Igarashi K. Angiographic findings of everolimus-eluting as compared to sirolimus-eluting stents: angiographic sub-study from the Randomized Evaluation of Sirolimus-eluting versus Everolimus-eluting stent Trial (RESET). *Cardiovasc Interv Ther.* 2013;28(4):344-51.
43. Khand A, Fisher M, Jones J, Patel B, Perry R, Mitsudo K. The collateral circulation of the heart in coronary total arterial occlusions in man: systematic review of assessment and pathophysiology. *Am Heart J.* 2013 Dec;166(6):941-52.
44. Toyota T, Furukawa Y, Ehara N, Funakoshi S, Morimoto T, Kaji S, Nakagawa Y, Kadota K, Iwabuchi M, Shiomi H, Yamamuro A, Kinoshita M, Kitai T, Kim K, Tani T, Kobori A, Kita T, Sakata R, Kimura T; Credo-Kyoto Investigators. Sex-based differences in clinical practice and outcomes for Japanese patients with acute myocardial infarction undergoing primary percutaneous coronary intervention. *Circ J.* 2013;77(6):1508-17.

45. Ozasa N, Morimoto T, Bao B, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Shiomi H, Tazaki J, Natsuaki M, Kimura T; CREDO-Kyoto Registry Investigators. β -blocker use in patients after percutaneous coronary interventions: one size fits all? Worse outcomes in patients without myocardial infarction or heart failure. *Int J Cardiol.* 2013;168(2):774-9.
46. Bao B, Ozasa N, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Shiomi H, Tada T, Tazaki J, Kato Y, Hayano M, Natsuaki M, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T, Kimura T. β -Blocker therapy and cardiovascular outcomes in patients who have undergone percutaneous coronary intervention after ST-elevation myocardial infarction. *Cardiovasc Interv Ther.* 2013;28(2):139-47.
47. Tokushige A, Shiomi H, Morimoto T, Ono K, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Hamasaki S, Tei C, Nakashima H, Mitsudo K, Nobuyoshi M, Kita T, Kimura T. Influence of initial acute myocardial infarction presentation on the outcome of surgical procedures after coronary stent implantation: a report from the CREDO-Kyoto PCI/CABG Registry Cohort-2. *Cardiovasc Interv Ther.* 2013;28(1):45-55.
48. Yamaji K, Kimura T, Morimoto T, Nakagawa Y, Inoue K, Nobuyoshi M, Mitsudo K; j-Cypher Registry Investigators. Percutaneous coronary intervention in patients with previous coronary artery bypass grafting (from the j-Cypher Registry). *Am J Cardiol.* 2013 Oct 15;112(8):1110-9.
49. Yamamoto E, Natsuaki M, Morimoto T, Furukawa Y, Nakagawa Y, Ono K, Mitsudo K, Nobuyoshi M, Doi O, Tamura T, Tanaka M, Kimura T; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Long-term outcomes after percutaneous coronary intervention for chronic total occlusion (from the CREDO-Kyoto registry cohort-2). *Am J Cardiol.* 2013 Sep 15;112(6):767-74.
50. Nishida K, Kimura T, Kawai K, Miyano I, Nakaoka Y, Yamamoto S, Kaname N, Seki S, Kubokawa S, Fukatani M, Hamashige N, Morimoto T, Mitsudo K; j-Cypher Registry Investigators. Comparison of outcomes using the sirolimus-eluting stent in calcified versus non-calcified native coronary lesions in patients on- versus not on-chronic hemodialysis (from the j-Cypher registry). *Am J Cardiol.* 2013 Sep 1;112(5):647-55.
51. Waki K, Arakaki Y, Mitsudo K. Long-term outcome of transcatheter

polytetrafluoroethylene-covered stent implantation in a giant coronary aneurysm of a child with Kawasaki disease. *Catheter Cardiovasc Interv.* 2013 Mar;81(4):713-6.

52. Kim MH, Yu LH, Tanaka H, Mitsudo K. Experience with a novel retrograde wiring technique for coronary chronic total occlusion. *J Interv Cardiol.* 2013 Jun;26(3):254-8.
53. Kozuma K, Kimura T, Suzuki N, Miyazawa A, Waseda K, Honda Y, Morimoto T, Aizawa T, Mitsudo K, Miyazaki S, Yamaguchi T, Isshiki T. Peri-stent contrast staining and very late stent thrombosis after sirolimus-eluting stent implantation: an observation from the RESTART (REgistry of Stent Thrombosis for review And Re-evaluaTion) angiographic substudy. *EuroIntervention.* 2013 Nov;9(7):831-40.
54. Kato M, Kimura T, Morimoto T, Nishikawa H, Uchida F, Suzuki H, Hayashi Y, Kadota K, Mitsudo K; J-Cypher Registry Investigators. Comparison of five-year outcome of sirolimus-eluting stent implantation for chronic total occlusions versus for non-chronic total occlusion (from the j-Cypher registry). *Am J Cardiol.* 2012;110(9):1282-9.
55. Masuno M, Watanabe A, Naing BT, Shimada T, Fujimoto W, Ninomiya S, Ueda Y, Kadota K, Kotaka T, Kondo E, Yamanouchi Y, Inoue M, Ouchi K, Kuroki Y. Ehlers-Danlos syndrome, vascular type: a novel missense mutation in the COL3A1 gene. *Congenit Anom (Kyoto).* 2012;52(4):207-10.
56. Kimura T, Morimoto T, Natsuaki M, Shiomi H, Igarashi K, Kadota K, Tanabe K, Morino Y, Akasaka T, Takatsu Y, Nishikawa H, Yamamoto Y, Nakagawa Y, Hayashi Y, Iwabuchi M, Umeda H, Kawai K, Okada H, Kimura K, Simonton CA, Kozuma K; RESET Investigators. Comparison of everolimus-eluting and sirolimus-eluting coronary stents: 1-year outcomes from the Randomized Evaluation of Sirolimus-eluting Versus Everolimus-eluting stent Trial (RESET). *Circulation.* 2012;126(10):1225-36.
57. Nakamura M, Otsuka Y, Ueda Y, Mitsudo K. Favorable pharmacokinetics of biolimus A9 after deployment of Nobori stent for coronary artery disease: insights from Nobori PK study in Japanese subjects. *Cardiovasc Interv Ther.* 2012 Jan;27(1):24-30.
58. Saito S, Nakamura S, Fujii K, Nakamura M, Isshiki T, Hirayama H, Kikuchi T, Fujita H, Nonogi H, Mitsudo K, Kimura T, Igarashi K, Saito K, Lansky AJ, Stone GW, Honda Y, Waseda K, Fitzgerald PJ, Sudhir K. Mid-term results of everolimus-eluting stent in a Japanese

population compared with a US randomized cohort: SPIRIT III Japan Registry with harmonization by doing. *J Invasive Cardiol.* 2012 Sep;24(9):444-50.

59. Shizuta S, Ando K, Nobuyoshi M, Ikeda T, Yoshino H, Hiramatsu S, Kazatani Y, Yamashiro K, Okajima K, Kajiya T, Kobayashi Y, Kato T, Fujii S, Mitsudo K, Inoue K, Ito H, Haruna Y, Doi T, Nishio Y, Ozasa N, Nishiyama K, Kita T, Morimoto T, Kimura T; PREVENT-SCD Investigators. Prognostic utility of T-wave alternans in a real-world population of patients with left ventricular dysfunction: the PREVENT-SCD study. *Clin Res Cardiol.* 2012 Feb;101(2):89-99.
60. Shiomi H, Morimoto T, Hayano M, Furukawa Y, Nakagawa Y, Tazaki J, Imai M, Yamaji K, Tada T, Natsuaki M, Saijo S, Funakoshi S, Nagao K, Hanazawa K, Ehara N, Kadota K, Iwabuchi M, Shizuta S, Abe M, Sakata R, Okabayashi H, Hanyu M, Yamazaki F, Shimamoto M, Nishiwaki N, Imoto Y, Komiya T, Horie M, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T, Kimura T; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Comparison of long-term outcome after percutaneous coronary intervention versus coronary artery bypass grafting in patients with unprotected left main coronary artery disease (from the CREDO-Kyoto PCI/CABG Registry Cohort-2). *Am J Cardiol.* 2012;110(7):924-32.
61. Kimura T, Morimoto T, Nakagawa Y, Kadota K, Nozaki Y, Tada T, Take S, Shiota K, Ito A, Nakashima H, Fujita H, Kawasaki T, Inada T, Nakao K, Miyazaki S, Doi O, Isshiki T, Nobuyoshi M, Mitsudo K. Antiplatelet therapy and long-term clinical outcome after sirolimus-eluting stent implantation: 5-year outcome of the j-Cypher registry. *Cardiovasc Interv Ther.* 2012;27(3):181-8.
62. Nakao T, Kimura T, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Nobuyoshi M, Kita T, Mitsudo K. The long-term efficacy of cilostazol in addition to dual antiplatelet therapy after sirolimus-eluting stent implantation for Japanese patients: an analysis of the 3-year follow-up outcomes from the j-Cypher registry. *Cardiovasc Interv Ther.* 2012;27(3):161-7.
63. Natsuaki M, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Shiomi H, Kimura T. Comparison of 3-year clinical outcomes after transradial versus transfemoral percutaneous coronary intervention. *Cardiovasc Interv Ther.* 2012;27(2):84-92.
64. Miyauchi K, Daida H, Morimoto T, Hiro T, Kimura T, Nakagawa Y, Yamagishi M, Ozaki Y, Kadota K, Kimura K, Hirayama A, Kimura K, Hasegawa Y, Uchiyama S, Matsuzaki M;

JAPAN-ACS Investigators. Reverse vessel remodeling but not coronary plaque regression could predict future cardiovascular events in ACS patients with intensive statin therapy--the extended JAPAN-ACS study. *Circ J.* 2012;76(4):825-32.

65. Tokushige A, Shiomi H, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Ehara N, Inada T, Kaburagi S, Hamasaki S, Tei C, Nakashima H, Ogawa H, Tatami R, Suwa S, Takizawa A, Nohara R, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T, Kimura T; CREDO-Kyoto PCI/CABG registry cohort-2 investigators. Incidence and outcome of surgical procedures after coronary bare-metal and drug-eluting stent implantation: a report from the CREDO-Kyoto PCI/CABG registry cohort-2. *Circ Cardiovasc Interv.* 2012;5(2):237-46.
66. Natsuaki M, Nakagawa Y, Morimoto T, Ono K, Shizuta S, Furukawa Y, Kadota K, Iwabuchi M, Kato Y, Suwa S, Inada T, Doi O, Takizawa A, Nobuyoshi M, Kita T, Kimura T; CREDO-Kyoto PCI/CABG Registry Cohort-2 Investigators. Impact of statin therapy on late target lesion revascularization after sirolimus-eluting stent implantation (from the CREDO-Kyoto Registry Cohort-2). *Am J Cardiol.* 2012;109(10):1387-96.
67. Toyofuku M, Kimura T, Morimoto T, Hayashi Y, Shiode N, Okimoto T, Otsuka M, Tamekiyo H, Tamura T, Kadota K, Inoue K, Mitsudo K; j-Cypher Registry investigators. Comparison of target-lesion revascularisation between left main coronary artery bifurcations and left anterior descending coronary artery bifurcations using the one and two stent approach with sirolimus-eluting stents. *EuroIntervention.* 2011;7(7):796-804.
68. Tada T, Kimura T, Morimoto T, Ono K, Furukawa Y, Nakagawa Y, Nakashima H, Ito A, Shiode N, Namura M, Inoue N, Nishikawa H, Nakao K, Mitsudo K; j-Cypher Registry Investigators. Comparison of three-year clinical outcomes after sirolimus-eluting stent implantation among insulin-treated diabetic, non-insulin-treated diabetic, and non diabetic patients from j-Cypher registry. *Am J Cardiol.* 2011 Apr 15;107(8):1155-62.
69. Kishi K, Kimura T, Morimoto T, Namura M, Muramatsu T, Nishikawa H, Hiasa Y, Isshiki T, Nobuyoshi M, Mitsudo K; j-Cypher Registry Investigators. Sirolimus-eluting stent implantation for ostial left anterior descending coronary artery lesions: three-year outcome from the j-Cypher Registry. *Circ Cardiovasc Interv.* 2011 Aug;4(4):362-70.
70. Tamura T, Kimura T, Morimoto T, Nakagawa Y, Furukawa Y, Kadota K, Tatami R, Kawai K,

- Sone T, Miyazaki S, Mitsudo K; j-Cypher Registry Investigators. Three-year outcome of sirolimus-eluting stent implantation in coronary bifurcation lesions: the provisional side branch stenting approach versus the elective two-stent approach. *EuroIntervention*. 2011;7(5):588-96.
71. Kimura T, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Shiomi H, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Tamura T, Shirotani M, Miki S, Matsuda M, Takahashi M, Ishii K, Tanaka M, Aoyama T, Doi O, Hattori R, Tatami R, Suwa S, Takizawa A, Takatsu Y, Takahashi M, Kato H, Takeda T, Lee JD, Nohara R, Ogawa H, Tei C, Horie M, Kambara H, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T. Long-term safety and efficacy of sirolimus-eluting stents versus bare-metal stents in real world clinical practice in Japan. *Cardiovasc Interv Ther*. 2011;26(3):234-45.
 72. Otsuka Y, Saito S, Nakamura M, Shuto H, Mitsudo K. Comparison of pharmacokinetics of the limus-eluting stents in Japanese patients. *Catheter Cardiovasc Interv*. 2011 Dec 1;78(7):1078-85.
 73. Kimura T, Morimoto T, Furukawa Y, Nakagawa Y, Kadota K, Iwabuchi M, Shizuta S, Shiomi H, Tada T, Tazaki J, Kato Y, Hayano M, Abe M, Tamura T, Shirotani M, Miki S, Matsuda M, Takahashi M, Ishii K, Tanaka M, Aoyama T, Doi O, Hattori R, Tatami R, Suwa S, Takizawa A, Takatsu Y, Takahashi M, Kato H, Takeda T, Lee JD, Nohara R, Tei C, Horie M, Kambara H, Fujiwara H, Mitsudo K, Nobuyoshi M, Kita T. Association of the use of proton pump inhibitors with adverse cardiovascular and bleeding outcomes after percutaneous coronary intervention in the Japanese real world clinical practice. *Cardiovasc Interv Ther*. 2011;26(3):222-33.
 74. Morino Y, Abe M, Morimoto T, Kimura T, Hayashi Y, Muramatsu T, Ochiai M, Noguchi Y, Kato K, Shibata Y, Hiasa Y, Doi O, Yamashita T, Hinohara T, Tanaka H, Mitsudo K; J-CTO Registry Investigators. Predicting successful guidewire crossing through chronic total occlusion of native coronary lesions within 30 minutes: the J-CTO (Multicenter CTO Registry in Japan) score as a difficulty grading and time assessment tool. *JACC Cardiovasc Interv*. 2011 Feb;4(2):213-21.
 75. Tada T, Kimura T, Morimoto T, Ono K, Furukawa Y, Nakagawa Y, Nakashima H, Ito A, Siode N, Namura M, Inoue N, Nishikawa H, Nakao K, Mitsudo K; j-Cypher Registry Investigators. Comparison of three-year clinical outcomes after sirolimus-eluting stent implantation among insulin-treated diabetic, non-insulin-treated diabetic, and non diabetic patients from j-Cypher registry. *Am J Cardiol*. 2011 Apr 15;107(8):1155-62.

76. Kawaguchi R, Kimura T, Morimoto T, Oshima S, Hoshizaki H, Kawai K, Shiode N, Hiasa Y, Mitsudo K; j-Cypher Registry Investigators. Safety and efficacy of sirolimus-eluting stent implantation in patients with acute coronary syndrome in the real world. *Am J Cardiol.* 2010 Dec 1;106(11):1550-60.
77. Nakamura M, Yokoi H, Hamazaki Y, Watarai M, Kijima M, Mitsudo K; Cypher J-PMS Investigators. Impact of insulin-treated diabetes and hemodialysis on long-term clinical outcomes following sirolimus-eluting stent deployment. Insights from a sub-study of the Cypher Stent Japan Post-Marketing Surveillance(Cypher J-PMS) Registry. *Circ J.* 2010 Nov;74(12):2592-7. Epub 2010 Oct 28.
78. Ozasa N, Kimura T, Morimoto T, Hou H, Tamura T, Shizuta S, Nakagawa Y, Furukawa Y, Hayashi Y, Nakao K, Matsuzaki M, Nobuyoshi M, Mitsudo K; j-Cypher Registry Investigators. Lack of effect of oral beta-blocker therapy at discharge on long-term clinical outcomes of ST-segment elevation acute myocardial infarction after primary percutaneous coronary intervention. *Am J Cardiol.* 2010 Nov 1;106(9):1225-33.
79. Kimura T, Morimoto T, Kozuma K, Honda Y, Kume T, Aizawa T, Mitsudo K, Miyazaki S, Yamaguchi T, Hiyoshi E, Nishimura E, Isshiki T; RESTART Investigators. Comparisons of baseline demographics, clinical presentation, and long-term outcome among patients with early, late, and very late stent thrombosis of sirolimus-eluting stents: Observations from the Registry of Stent Thrombosis for Review and Reevaluation (RESTART). *Circulation.* 2010 Jul 6;122(1):52-61.
80. Abe M, Kimura T, Morimoto T, Taniguchi T, Yamanaka F, Nakao K, Yagi N, Kokubu N, Kasahara Y, Kataoka Y, Otsuka Y, Kawamura A, Miyazaki S, Nakao K, Horiuchi K, Ito A, Hoshizaki H, Kawaguchi R, Setoguchi M, Inada T, Kishi K, Sakamoto H, Morioka N, Imai M, Shiomi H, Nonogi H, Mitsudo K; j-Cypher Registry Investigators. Sirolimus eluting stent versus balloon angioplasty for sirolimus-eluting stent restenosis: Insights from the j-Cypher Registry. *Circulation.* 2010 Jul 6;122(1):42-51.
81. Shirai S, Kimura T, Nobuyoshi M, Morimoto T, Ando K, Soga Y, Yamaji K, Kondo K, Sakai K, Arita T, Goya M, Iwabuchi M, Yokoi H, Nosaka H, Mitsudo K; j-Cypher Registry Investigators. Impact of multiple and long sirolimus-eluting stent implantation on 3-year clinical outcomes in the j-Cypher Registry. *JACC Cardiovasc Interv.* 2010 Feb;3(2):180-8.

82. Morino Y, Kimura T, Hayashi Y, Muramatsu T, Ochiai M, Noguchi Y, Kato K, Shibata Y, Hiasa Y, Doi O, Yamashita T, Morimoto T, Abe M, Hinohara T, Mitsudo K; J-CTO Registry Investigators. In-hospital outcomes of contemporary percutaneous coronary intervention in patients with chronic total occlusion insights from the J-CTO Registry (Multicenter CTO Registry in Japan). *JACC Cardiovasc Interv.* 2010 Feb;3(2):143-51.
83. Kimura T, Isshiki T, Hayashi Y, Oshima S, Namura M, Nakashima H, Kawai K, Sone T, Tatami R, Meguro T, Nobuyoshi M, Mitsudo K. Incidence and outcome of surgical procedures after sirolimus-eluting stent implantation: a report from the j-Cypher registry. *Cardiovasc Interv Ther.* 2010 Jan;25(1):29-39.
84. Kim MH, Yu LH, Mitsudo K. A new retrograde wiring technique for chronic total occlusion. *Catheter Cardiovasc Interv.* 2010 Jan 1;75(1):117-9.
85. Kimura T, Morimoto T, Nakagawa Y, Tamura T, Kadota K, Yasumoto H, Nishikawa H, Hiasa Y, Muramatsu T, Meguro T, Inoue N, Honda H, Hayashi Y, Miyazaki S, Oshima S, Honda T, Shiode N, Namura M, Sone T, Nobuyoshi M, Kita T, Mitsudo K; j-Cypher Registry Investigators. Antiplatelet therapy and stent thrombosis after sirolimus-eluting stent implantation. *Circulation.* 2009;119(7):987-95.
86. Toyofuku M, Kimura T, Morimoto T, Hayashi Y, Ueda H, Kawai K, Nozaki Y, Hiramatsu S, Miura A, Yokoi Y, Toyoshima S, Nakashima H, Haze K, Tanaka M, Take S, Saito S, Isshiki T, Mitsudo K; j-Cypher Registry Investigators. Three year outcomes after sirolimus-eluting stent implantation for unprotected left main coronary artery disease: insights from the j-Cypher registry. *Circulation.* 2009 Nov 10;120(19):1866-74.
87. Yoshida K, Endo H, Sadamasa N, Narumi O, Chin M, Inoue K, Mitsudo K, Yamagata S. Evaluation of carotid artery atherosclerotic plaque distribution by using long-axis high-resolution black-blood magnetic resonance imaging. *J Neurosurg.* 2008 Dec;109(6):1042-8.
88. Inoue K, Mitsudo K, Nobuyoshi M. Medial necrosis due to sirolimus-eluting stent implantation in human coronary artery. *J Cardiol.* 2008 Feb;51(1):60-4.
89. Morino Y, Yamamoto H, Mitsudo K, Nagaoka M, Takeuchi H, Okamoto N, Kozuma K,

Matsuzaki A, Tanabe K, Hara K, Tanabe T, Ikari Y. Functional formula to determine adequate balloon diameter of simultaneous kissing balloon technique for treatment of bifurcated coronary lesions: clinical validation by volumetric intravascular ultrasound analysis. *Circ J.* 2008 Jun;72(6):886-92.

90. Inoue K, Kawahara K, Biswas KK, Ando K, Mitsudo K, Nobuyoshi M, Maruyama I. HMGB1 expression by activated vascular smooth muscle cells in advanced human atherosclerosis plaques. *Cardiovasc Pathol.* 2007 May-Jun;16(3):136-43.
91. Shinozaki N, Yokoi H, Iwabuchi M, Nosaka H, Kadota K, Mitsudo K, Nobuyoshi M. Initial and follow-up results of the BiodivYsio phosphorylcholine coated stent for treatment of coronary artery disease. *Circ J.* 2005;69(3):295-300.
92. Yokoi H, Nobuyoshi M, Mitsudo K, Kawaguchi A, Yamamoto A; ATHEROMA Study Investigators. Three-year follow-up results of angiographic intervention trial using an HMG-CoA reductase inhibitor to evaluate retardation of obstructive multiple atheroma (ATHEROMA) study. *Circ J.* 2005 Aug;69(8):875-83.
93. Stone GW, Reifart NJ, Moussa I, Hoyer A, Cox DA, Colombo A, Baim DS, Teirstein PS, Strauss BH, Selmon M, Mintz GS, Katoh O, Mitsudo K, Suzuki T, Tamai H, Grube E, Cannon LA, Kandzari DE, Reisman M, Schwartz RS, Bailey S, Dangas G, Mehran R, Abizaid A, Moses JW, Leon MB, Serruys PW. Percutaneous recanalization of chronically occluded coronary arteries: a consensus document: part II. *Circulation.* 2005 Oct 18;112(16):2530-7.
94. Stone GW, Kandzari DE, Mehran R, Colombo A, Schwartz RS, Bailey S, Moussa I, Teirstein PS, Dangas G, Baim DS, Selmon M, Strauss BH, Tamai H, Suzuki T, Mitsudo K, Katoh O, Cox DA, Hoyer A, Mintz GS, Grube E, Cannon LA, Reifart NJ, Reisman M, Abizaid A, Moses JW, Leon MB, Serruys PW. Percutaneous recanalization of chronically occluded coronary arteries: a consensus document: part I. *Circulation.* 2005 Oct 11;112(15):2364-72.
95. Stone GW, Colombo A, Teirstein PS, Moses JW, Leon MB, Reifart NJ, Mintz GS, Hoyer A, Cox DA, Baim DS, Strauss BH, Selmon M, Moussa I, Suzuki T, Tamai H, Katoh O, Mitsudo K, Grube E, Cannon LA, Kandzari DE, Reisman M, Schwartz RS, Bailey S, Dangas G, Mehran R, Abizaid A, Serruys PW. Percutaneous recanalization of chronically occluded coronary arteries: procedural techniques, devices, and results. *Catheter Cardiovasc Interv.* 2005 Oct;66(2):217-36.

